Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MARCH 3.

Dr. GEO. H. HORN in the chair.

Twenty-two persons present.

MARCH 10.

The President, Dr. LEIDY, in the chair.

Twenty-six persons present.

Papers under the following titles were presented for publication :—

Fossil Faunas of Central Iowa. By Charles R. Keyes.

Notice of some Entozoa. By Joseph Leidy, M. D.

On the Young of Baculites compressus Say.—Mr. AMOS P. BROWN described the young of *Baculites compressus* Say, recently discovered by him in some cretaceous marl from the vicinity of Deadwood, South Dakota. Associated with them in the same material were several species of *Baculites, Scaphites* and *Inoceramus*. The young *Baculites* were of the form shown in figures 1 and 2 and varied in length from 1 to 3 cm., with a diameter of 0·4 to 2 mm. Other larger fragments with the spiral end broken off were found from 1·5 to 6 cm. in diameter. An examination of the form of the septa and suture lines showed the forms to belong to the Ammonitidae, and by the examination of an extensive series it was possible to determine the genus and species.